

# FUNCTIONS TILING WITH SEVERAL LATTICES

MIHAIL N. KOLOUNTZAKIS AND EFFIE PAPAGEORGIU

ABSTRACT. We study the problem of finding a function  $f$  with “small support” that simultaneously tiles with finitely many lattices  $\Lambda_1, \dots, \Lambda_N$  in  $d$ -dimensional Euclidean spaces. We prove several results, both upper bounds (constructions) and lower bounds on how large this support can and must be. We also study the problem in the setting of finite abelian groups, which turns out to be the most concrete setting. Several open questions are posed.

## CONTENTS

1. Introduction to the Steinhaus tiling problem and its variants	1
1.1. Previous results	2
1.2. Structure of this paper	2
2. The diameter and volume of soft multi-lattice tiles	3
2.1. Convolution tiles and their diameter	3
2.2. Common tiles whose support has small volume	6
2.3. Allowing for lattices with many relations	7
3. The common fundamental domain problem without measurability	10
4. The problem in finite abelian groups	11
References	15

## 1. INTRODUCTION TO THE STEINHAUS TILING PROBLEM AND ITS VARIANTS

The classical Steinhaus tiling problem concerns tiling by translations only. This is the version of tiling where we have a set (the tile), say in the plane, and we are *translating* it around in such a way that every point of the plane is covered exactly once by these translates. In the Steinhaus tiling problem we are seeking a tile that can tile *simultaneously* with many different sets of translations. The most important case is: can we find a subset of the plane which can tile (by translations) with all rotates of the integer lattice  $\mathbb{Z}^2$ ?

There are two major variations of the Steinhaus problem: the *measurable* and the *set-theoretic* case. In the measurable case we demand our tile to be a Lebesgue measurable subset of  $\mathbb{R}^d$  and we are, at the same time, relaxing our requirements and are allowing a subset of measure 0 of space not to be covered exactly once by the translates of the tile. In the set-theoretic case we allow the tile to be any subset and we typically ask that every point is covered exactly once, allowing no exceptions.

In this paper we are dealing with the measurable Steinhaus tiling problem. Here the tile is not necessarily a set but can be a measurable function. Tiling

---

*Date:* June 22, 2021.

*2020 Mathematics Subject Classification.* primary 52C22; secondary 20K99.

*Key words and phrases.* Lattices, tiling, common fundamental domain, Steinhaus problem.

Supported by the Hellenic Foundation for Research and Innovation, Project HFRI-FM17-1733 and by grant No 4725 of the University of Crete.

now means that this function  $f : \mathbb{R}^d \rightarrow \mathbb{R}$  is translated around by a countable set of translates  $T \subseteq \mathbb{R}^d$  and these copies add up to a constant, almost everywhere

$$\sum_{t \in T} f(x - t) = \text{const.}, \quad \text{for almost every } x \in \mathbb{R}^d.$$

Clearly tiling by a function is a generalization of tiling by a set.

Satisfying the requirements of the Steinhaus tiling problem with a function instead of with a set is generally much easier. The problem becomes interesting only if one asks for further properties that this function should have. Therefore we strive to find a function with *small support*, or to prove that the support of such a function must necessarily be large. Asking for  $f$  to have a small support goes against  $f$  having the ability to tile space, especially with many different sets of translations  $T$ . The reason is that for  $f$  to tile by translations with  $T$  its Fourier transform must contain a rich set of zeros [19]. This set of zeros must be able to support the Fourier transform of the measure  $\delta_T = \sum_{t \in T} \delta_t$  (which encodes the set of translations). By the well known uncertainty principle in harmonic analysis a rich set of zeros for  $\widehat{f}$  usually requires (in various different senses) a large support for  $f$  [12].

**1.1. Previous results.** Komjáth [23] answered the Steinhaus question in the affirmative in  $\mathbb{R}^2$  when tiling by all rotates of the set  $B = \mathbb{Z} \times \{0\}$  showing that there are such Steinhaus sets (but such a set  $A$  cannot be measurable as was shown recently in [21]). Sierpinski [28] showed that a bounded set  $A$  which is either closed or open cannot have the lattice Steinhaus property (that is, intersect all rigid motions of  $\mathbb{Z}^2$  at exactly one point – another way to say that  $A$  tiles precisely with all rotates of  $\mathbb{Z}^2$ ). Croft [7] and Beck [1] showed that no bounded and measurable set  $A$  can have the lattice Steinhaus property (but see also [24]). Kolountzakis [17, 16] and Kolountzakis and Wolff [22] proved that any measurable set in the plane that has the measurable Steinhaus property must necessarily have very slow decay at infinity (any such set must have measure 1). In [22] it was also shown that there can be no measurable Steinhaus sets in dimension  $d \geq 3$  (tiling with all rotates  $\rho\mathbb{Z}^d$ , where  $\rho$  is in the full orthogonal group) a fact that was also shown later by Kolountzakis and Papadimitrakis [20] by a very different method. See also [5, 25, 6, 29]. Kolountzakis [18] looks at the case where we are only asking for our set to tile with *finitely many* lattices, not all rotates as in the original problem, which we are also doing in this paper. In a major result Jackson and Mauldin [14, 13] proved the existence of Steinhaus sets in the plane which tile with all rotates of  $\mathbb{Z}^2$  (not necessarily measurable). Their method does not extend to higher dimension  $d \geq 3$ . See also [26, 15]. It was also shown in [21] that a set  $A$  which tiles with all rotates of a finite set  $B$  cannot be measurable.

**1.2. Structure of this paper.** In §2 we study the common tiles  $f$  for a finite collection of  $N$  lattices in  $\mathbb{R}^d$ . The emphasis is on the dependence of the size of the common tile on  $N$ .

In §2.1 we study the diameter of common tiles, a study that had begun in [22]. We show in Theorem 2.3 that there are cases of lattices for which we have a linear lower bound (matching the linear upper bound arising from convolution tiles) for the diameter of the support of any common tile.

In §2.2 we study the volume of the common tiles instead of their diameter. In Theorem 2.4 we show that convolution tiles with nonnegative convolution factors necessarily have volume  $\gtrsim N^d$  for their support. In Theorem 2.5 we give a simple non-trivial lower bound for the length for a convolution tile of two factors.

Then in §2.3 we turn away from lattices satisfying genericity conditions (e.g. having a direct sum) to lattices which satisfy many algebraic relations. We show that this helps greatly with the diameter and volume of the support of their common tiles (construction in Theorem 2.8 for  $d \geq 2$  and in Theorem 2.9 for  $d = 1$ ).

In §3 we study common tiles of lattices without the measurability assumption. For two lattices of the same volume in  $\mathbb{R}^d$  these were known to always exist [18]. We show in Theorem 3.3 that the condition of equal volume is actually necessary for the existence of a common fundamental domain, which had remained an open question in [18].

In §4 we study the problem of finding a common tiling function for subgroups of a finite abelian group, trying now to minimize the size (cardinality) of its support.

In Theorem 4.2 we show that we can reduce the problem to the case where the two subgroups have no intersection, i.e. we reduce the study to direct products of groups. Theorem 4.3 solves the problem exactly in the special case when the size of one group divides the size of the other. We then connect the problem to the problem of the support of *copulas* (a generalization of doubly stochastic matrices to the non-square case) as they have been studied in combinatorics and statistics and we deal with some special cases in Lemma 4.5. Finally in Theorem 4.6 we exploit this connection to produce, in some special cases again, a common tile for two lattices in  $\mathbb{R}$  which is slightly better in terms of its length than the convolution common tile.

Several open questions are posed throughout.

## 2. THE DIAMETER AND VOLUME OF SOFT MULTI-LATTICE TILES

**2.1. Convolution tiles and their diameter.** It has long been known [1, 17] that for a function  $f \in L^1(\mathbb{R}^d)$  to tile with all rotates of  $\mathbb{Z}^d$  it is necessary and sufficient that  $\widehat{f}$  vanishes on all spheres centered at the origin that contain any integer lattice point. This easily implies that for  $d \geq 2$  any such function must have unbounded support and even more quantitative lower bounds for the rate of decay of  $f$  near infinity [17, 16, 22].

This is no longer true if one restricts the number of rotates of  $\mathbb{Z}^d$  that we demand  $f$  tiles with. It makes sense to generalize the question and ask for a function  $f$  which tiles with a given set of lattices  $\Lambda_i \subseteq \mathbb{R}^d$ ,  $i = 1, 2, \dots, N$ , and such that

$$\text{diam supp } f$$

is small. We remind [22] that for  $f$  to tile with the lattices  $\Lambda_i$  it is necessary and sufficient for  $\widehat{f}$  to vanish on the dual lattices  $\Lambda_i^*$ , except at 0. (The dual lattice of a lattice  $\Lambda = A\mathbb{Z}^d$ , with  $A$  being a non-singular  $d \times d$  matrix, is the lattice  $\Lambda^* = A^{-\top}\mathbb{Z}^d$ .)

Therefore we have to find a function  $f$ , with support of small diameter, such that  $\widehat{f}$  is 0 on each  $\Lambda_i^* \setminus \{0\}$ . The first thing that comes to mind is to take  $f$  to be a convolution. It takes a moment to verify that if  $f$  tiles with a set of translates  $T$  then so does  $g * f$  for any  $g \in L^1(\mathbb{R}^d)$ . One can either verify this by checking the definition of tiling for  $g * f$  or observe that tiling is a condition that can be checked on the Fourier side [19] and  $\widehat{g * f} = \widehat{g} \cdot \widehat{f}$  has an even richer set of zeros than  $\widehat{f}$ .

So, since  $\widehat{f}$  has to vanish on the dual lattices  $\Lambda_i^* \setminus \{0\}$  we can take

$$(2.1) \quad f = \mathbb{1}_{D_1} * \mathbb{1}_{D_2} * \cdots * \mathbb{1}_{D_N},$$

where  $D_i$  is a fundamental parallelepiped of  $\Lambda_i$ . Since  $D_i + \Lambda_i$  is a tiling it follows that  $\widehat{\mathbb{1}_{D_i}}$  vanishes on  $\Lambda_i^* \setminus \{0\}$  and that  $f$  vanishes on their union and hence tiles with all  $\Lambda_i$ . This can be slightly generalized by taking, instead of the indicator functions  $\mathbb{1}_{D_i}$  any function  $f_i$  that tiles with  $\Lambda_i$

$$(2.2) \quad f = f_1 * f_2 * \cdots * f_N.$$

The following observation was already made in [22] in the case  $f_i = \mathbb{1}_{D_i}$ .

**Theorem 2.1.** *If  $\Lambda_1, \dots, \Lambda_N$  are lattices in  $\mathbb{R}^d$  of volume  $c_1 \leq \text{vol } \Lambda_i$  and  $f = f_1 * f_2 * \cdots * f_N$  then*

$$(2.3) \quad \text{diam supp } f \geq C_d N.$$

*Proof.* By the pigeonhole principle one can pick a coordinate axis, say the first one, such that at least  $N/d$  of the functions  $f_i$  have  $\text{supp } f_i$  project onto the first coordinate axis on a set of diameter  $\geq a_d$ , where  $a_d$  is a constant that depends on  $d$  only. For if not then we would be able to find a  $f_i$  whose support is contained in a cube of arbitrarily small side. Since  $f_i$  tiles with  $\Lambda_i$  its support has to contain at least one element from almost all of the cosets of  $\Lambda_i$  in  $\mathbb{R}^d$ . This contradicts our assumed lower bound on  $\text{vol } \Lambda_i$ .

The Titchmarsh convolution theorem [8] says that

$$\text{co supp } (A + B) = \text{co supp } A + \text{co supp } B,$$

where  $\text{co}$  denotes the convex hull of a set and  $A, B$  are two arbitrary integrable functions of compact support.

If we write  $\phi$  for the convolution of those  $f_i$  that we collected in the first paragraph of this proof and  $\psi$  for the remaining  $f_i$  then  $f = \phi * \psi$  and

$$\text{co supp } f = \text{co supp } \phi + \text{co supp } \psi,$$

which implies that

$$(2.4) \quad \text{diam supp } f = \text{diam co supp } f \geq \text{diam co supp } \phi \geq \text{diam } \pi_1 \text{co supp } \phi,$$

where  $\pi_1$  denotes projection onto the first coordinate axis.

Again by the Titchmarsh convolution theorem,  $\pi_1 \text{co supp } \phi$  is the sum of the  $\pi_1 \text{co supp } f_i$  for those  $f_i$  that participate in the definition of  $\phi$  and for these we know that

$$\text{diam } \pi_1 \text{co supp } f_i \geq \text{diam } \pi_1 \text{supp } f_i \geq a_d.$$

But for any two one-dimensional sets  $E, F \subseteq \mathbb{R}$  we have  $\text{diam } (E + F) = \text{diam } E + \text{diam } F$ , which implies, using (2.4), that

$$\text{diam supp } f \geq \frac{a_d}{d} N,$$

as we had to prove. □

If the lattices  $\Lambda_i$  satisfy some “roundness” assumption, e.g. if each  $\Lambda_i$  is assumed to have a fundamental domain of diameter bounded independent of  $N$  (as in the important case when all the lattices are rotates of  $\mathbb{Z}^d$ ), then the convolution tile (2.1) clearly has diameter which is also at most  $C \cdot N$ .

On the other hand we have the following rather general lower bound for the diameter of the support [22] assuming only a certain genericity assumption (2.5) on the  $\Lambda_i$ .

**Theorem 2.2.** *If  $\Lambda_1, \dots, \Lambda_N \subseteq \mathbb{R}^d$ ,  $d \geq 1$ , are lattices of volume equal to 1 such that*

$$(2.5) \quad \Lambda_i \cap \Lambda_j = \{0\} \quad \text{for all } i \neq j,$$

*then if  $f$  tiles with all these lattices we have*

$$(2.6) \quad \text{diam supp } f \geq C_d N^{1/d}.$$

The main question is therefore:

**Question 1.** *Can the gap between the lower bound (2.6) and the linear upper bound  $O(N)$  achievable by the convolution tile (2.1) (in the case of “round” lattices, having fundamental domains bounded in diameter by a constant) be bridged?*

*Are there examples of lattices  $\Lambda_i$ ,  $i = 1, 2, \dots, N$ , satisfying (2.5) and a non-zero function  $f \in L^1(\mathbb{R}^d)$  that tiles with all  $\Lambda_i$  and such that*

$$\text{diam supp } f = o(N)?$$

*In other words, do there exist collections of lattices for which a common tile  $f$  can be found which is diameter-wise more efficient than the convolution construction (2.1)?*

Our next result is that for some collections of lattices the linear upper bound cannot be improved. The lattices given are both “round” (have a fundamental domain bounded independent of  $N$ ) and satisfy the genericity assumption (2.5). There are however collinearities so, in some sense, this is not a generic situation.

**Theorem 2.3.** *For  $d \geq 1$  and for each  $N$  there are lattices  $\Lambda_1, \dots, \Lambda_N \subseteq \mathbb{R}^d$ , of volume 1, such that if  $f \in L^1(\mathbb{R}^d)$ ,  $\int f \neq 0$ , tiles with all of them then*

$$\text{diam supp } f \geq C_d N.$$

*Proof.* We give the proof in the case  $d = 2$ . It works with obvious changes in all dimensions  $d > 2$  and it is even easier in dimension  $d = 1$ .

Take  $\Lambda_i^*$  to be generated by the two vectors

$$u_i = (0, a_i), v_i = (1/a_i, 0),$$

where the numbers  $a_1, \dots, a_N$  are linearly independent over  $\mathbb{Q}$  and

$$0.9 < a_i < 1.$$

If  $f$  tiles with all  $\Lambda_i$  then  $\widehat{f}$  vanishes on all points of the form

$$(0, k \cdot a_i), \quad i = 1, 2, \dots, N, \quad k \in \mathbb{Z} \setminus \{0\}.$$

Since all these points are different it follows that the density of zeros on the  $y$ -axis is  $\geq C \cdot N$ . This implies that

$$\text{diam supp } \pi_2(f) \geq C \cdot N$$

(say, by Jensen's formula) where  $\pi_2(f)$  is the one-variable function

$$\pi_2(f)(y) = \int_{\mathbb{R}} f(x, y) dx.$$

(This is not an identically zero function by our assumption on the integral of  $f$ .) This in turn implies

$$\text{diam supp } f \geq C \cdot N.$$

□

**2.2. Common tiles whose support has small volume.** Another measure of smallness of the support is its volume. Can we construct a common tile  $f$  for the lattices  $\Lambda_i$  such that  $|\text{supp } f|$  is small?

In the case of  $f$  given by (2.1) it is clear that

$$\text{supp } f = D_1 + D_2 + \dots + D_N.$$

To keep things concrete let us assume that all  $|D_i| = 1$  in (2.1) (unimodular lattices). Then the Brunn-Minkowski inequality [10] says that

$$|\text{supp } f| = |D_1 + \dots + D_N| \geq (|D_1|^{1/d} + \dots + |D_N|^{1/d})^d \geq N^d.$$

This lower bound

$$|\text{supp } f| \geq CN^d$$

clearly holds also for functions of the form

$$(2.7) \quad f = f_1 * f_2 * \dots * f_N, \quad f_i \geq 0,$$

where for all  $i = 1, 2, \dots, N$  we assume that the *nonnegative* function  $f$  tiles with  $\Lambda_i$ .

We have proved:

**Theorem 2.4.** *For any collection of lattices  $\Lambda_1, \dots, \Lambda_N$  in  $\mathbb{R}^d$  of volume at least 1 and any common tile  $f$  for them of the form*

$$f = f_1 * f_2 * \dots * f_N, \quad f_i \geq 0,$$

*with  $f_i$  tiling with  $\Lambda_i$ , we have*

$$|\text{supp } f| \geq N^d.$$

But when the functions  $f$  are signed (or complex) we only have

$$\text{supp } f \subseteq \text{supp } f_1 + \dots + \text{supp } f_N,$$

not necessarily equality, which brings us to the next question.

**Question 2.** *If  $f$  is given by (2.7), is it true that*

$$(2.8) \quad |\text{supp } f| \geq CN^d?$$

If one requires that the lattices  $\Lambda_1, \Lambda_2, \dots, \Lambda_N \subset \mathbb{R}^d$  have the same volume, say 1, and the sum  $\Lambda_1^* + \Lambda_2^* + \dots + \Lambda_N^*$  of their dual lattices is direct, then, by [18, Theorem 2], they possess a measurable common almost fundamental domain  $E$  (generally unbounded). In this case,  $|E| = \text{vol}(\Lambda_i) = 1$ . So then one can take  $f = \mathbb{1}_E$ , which tiles with all  $\Lambda_i$ ,  $i = 1, 2, \dots, N$ , with  $|\text{supp } f| = |E| = 1$ .

Motivated by the previous observation, but now dropping the equal volume assumption, we ask the following:

**Question 3.** Consider the lattices  $\Lambda_1, \Lambda_2, \dots, \Lambda_N$ , with  $\frac{1}{2} \leq \text{vol}(\Lambda_i) \leq 2$ . Is there a function  $f$  that tiles with all  $\Lambda_i$ , such that

$$|\text{supp } f| = o(N^d)?$$

In the simplest case in dimension  $d = 1$ , and for two lattices only, a basic question seems to be to ask if the convolution (2.7) is best in terms of the length of the support. Here we can offer a simple lower bound assuming a nonnegative function.

**Theorem 2.5.** Suppose the nonnegative  $f : \mathbb{R} \rightarrow \mathbb{R}^{\geq 0}$  is measurable and tiles with both  $\Lambda_1 = \mathbb{Z}$  and with  $\Lambda_2 = \alpha\mathbb{Z}$ , where  $\alpha \in (0, 1)$ :

$$(2.9) \quad \sum_{n \in \mathbb{Z}} f(x - n) = 1, \quad \sum_{n \in \mathbb{Z}} f(x - n\alpha) = \frac{1}{\alpha}, \quad \text{for almost every } x \in \mathbb{R}.$$

Then

$$(2.10) \quad |\text{supp } f| \geq \left\lceil \frac{1}{\alpha} \right\rceil \alpha \geq 2\alpha.$$

*Remark 2.6.* If we assume the first equation in (2.9) then the constant in the second equation is forced to be  $1/\alpha$ . This is because  $\int f = 1$  (from the first equation), so repeating  $f$  at a set of translates of density  $1/\alpha$  will give a constant (assuming it tiles) at that level.

*Remark 2.7.* Notice that if  $\alpha$  is just a little less than 1 then (2.10) gives a lower bound of  $2\alpha$ , which shows that the convolution  $\mathbb{1}_{[0,1]} * \mathbb{1}_{[0,\alpha]}$  is almost optimal in this case, having support of size  $1 + \alpha$ .

But if, on the other hand,  $\alpha$  is just over  $1/2$  then the lower bound is just over 1 but the convolution upper bound is just over  $3/2$ , a considerable gap.

*Proof.* From the first equation in (2.9) it follows that  $f(x) \leq 1$  for almost every  $x$ . For the second equation to be true it therefore follows that for almost every  $x \in \mathbb{R}$  there are at least  $\lceil 1/\alpha \rceil$  different values of  $n \in \mathbb{Z}$  such that  $f(x - n\alpha) > 0$ . Using this for almost all  $x \in [0, \alpha)$  (which ensures that for different  $x$  the locations  $x - n\alpha$  are also different) gives (2.10).  $\square$

**Question 4.** What is the least possible length of the support of  $f$  for a nonnegative  $f$  that tiles with both  $\mathbb{Z}$  and  $\alpha\mathbb{Z}$ ? See Remark 4.7 for a special case.

**2.3. Allowing for lattices with many relations.** If we have  $N$  lattices

$$\Lambda_1, \dots, \Lambda_N \subseteq \mathbb{R}^d$$

we can find a function that tiles with them all, namely the function  $f$  in (2.1). If our lattices are assumed to each have a fundamental domain bounded by  $\sim 1$  then  $\text{diam supp } f = O(N)$ , and this cannot be improved for functions  $f$  arising from (2.1). We show here that we can choose the lattices  $\Lambda_j$  so that a common tiling function exists which is much more tight than that, tighter

even than what Theorem 2.2 imposes. Of course our lattices will not satisfy the genericity condition (2.5) of Theorem 2.2, but will satisfy a lot of relations (their intersection will be a large lattice, in terms of density).

Fix a large prime  $p$  and consider the group  $\mathbb{Z}_p^d$ . Any nonzero element  $g$  of this group generates a cyclic subgroup of order  $p$ . It follows that  $\mathbb{Z}_p^d$  has

$$\frac{p^d - 1}{p - 1} \sim p^{d-1} =: N$$

different cyclic subgroups. For each such subgroup  $G$ , which we now view as a subset of  $\{0, 1, \dots, p - 1\}^d$ , consider the lattice

$$\Lambda_G = (p\mathbb{Z})^d + G,$$

which contains the lattice  $\Lambda = (p\mathbb{Z})^d$  and has volume

$$\text{vol } \Lambda_G = \frac{\text{vol } (p\mathbb{Z})^d}{|G|} = p^{d-1} = N.$$

The function  $f = \mathbb{1}_{[0,p)^d}$ ,  $[0,p)^d$  being a fundamental domain of  $\Lambda$ , tiles with  $\Lambda$  and, therefore, with any larger group, so  $f$  is a common tile of all  $\Lambda_G$ .

In order to make the volume of the  $\Lambda_G$  equal to 1 we shrink everything by  $N^{1/d}$ :

$$\Lambda'_G = N^{-1/d} \Lambda_G, \quad f'(x) = f(N^{1/d}x).$$

So we have  $\sim N$  lattices  $\Lambda'_G$  of volume 1 and a common tile  $f'$  for them with

$$\text{diam supp } f' = \text{diam supp } f \cdot N^{-1/d} = \sqrt{d} p N^{-1/d} = \sqrt{d} N^{\frac{1}{d-1} - \frac{1}{d}} = \sqrt{d} N^{\frac{1}{d(d-1)}}.$$

We have proved:

**Theorem 2.8.** *In dimension  $d \geq 2$  and for arbitrarily large  $N$  we can find  $N$  lattices of volume 1 and a common tile  $f$  for them with*

$$\text{diam supp } f = O_d \left( N^{\frac{1}{d(d-1)}} \right),$$

and, consequently, with

$$|\text{supp } f| = O_d \left( N^{\frac{1}{d-1}} \right).$$

**Question 5.** *Derive a lower bound for  $\text{diam supp } f$ , for  $f$  tiling with  $\Lambda_1, \dots, \Lambda_N \subseteq \mathbb{R}^d$  and with  $f \geq 0$  (or just  $\int f > 0$ ) under no algebraic conditions for the lattices  $\Lambda_j$ , assuming only that  $\text{vol } \Lambda_j \sim 1$ .*

The construction that we used to prove Theorem 2.8 gives nothing in dimension  $d = 1$ . Yet, we can prove that, if we allow relations among the lattices, we can achieve  $\text{diam supp } f = o(N)$  in dimension 1 as well.

Let us start by defining

$$\lambda_j = \frac{1}{N + j}, \quad \Lambda_j = \lambda_j \mathbb{Z}, \quad (j = 1, 2, \dots, N).$$

We will first construct a function  $f$  which tiles with all the  $\Lambda_j$ ,  $j = 1, 2, \dots, N$ , such that

$$\text{diam supp } f = o(1).$$

The Fourier transform of such an  $f$  must vanish on the dual lattices

$$\Lambda_j^* = \lambda_j^{-1} \mathbb{Z} = (N + j) \mathbb{Z}, \quad (j = 1, 2, \dots, N)$$

except at 0. Write

$$U = \bigcup_{j=1}^N (N+j)\mathbb{Z} \setminus \{0\}.$$

By a result of Erdős [9]  $U$ , the set of integers which are divisible by one of the integers in  $\{N+1, N+2, \dots, 2N\}$ , has density tending to 0 with  $N$ . Tenenbaum [30] has given the estimate that this density is at most

$$(2.11) \quad \frac{1}{\log^{\delta-o(1)} N'}$$

where  $\delta = 0.086071 \dots$  is an explicit constant.

It is an important result of Beurling [2] that if  $\Lambda$  is a uniformly discrete set of real numbers of upper density  $\rho$  then for any  $\epsilon > 0$  we can find a continuous function  $f$ , not identically zero, supported by the interval  $[0, \rho + \epsilon]$  such that  $\widehat{f}(\lambda) = 0$  for all  $\lambda \in \Lambda$ . We can even ask that  $\widehat{f}(0) = 1$  if  $0 \notin \Lambda$ . By Tenenbaum's estimate (2.11) we can take  $\rho = \log^{-\delta+o(1)} N$  and the set  $U$ , being a set of integers and thus uniformly discrete, satisfies the assumptions of Beurling's theorem, so there is a function  $f$  supported in the interval  $[0, \log^{-\delta+o(1)} N]$ , with integral 1, such that  $\widehat{f} = 0$  on  $U$ . It follows that  $f$  tiles with all  $\Lambda_j$ .

We now scale by a factor of  $N$

$$f'(x) = f(x/N), \quad \Lambda'_j = N\Lambda_j, \quad \text{diam supp } f' = O(N \log^{-\delta+o(1)} N)$$

and obtain the first half of the following theorem.

**Theorem 2.9.** *We can find  $N$  lattices  $\Lambda_j \subseteq \mathbb{R}$  of with  $\text{vol } \Lambda_j \sim 1$  and a function  $f$  with  $\int f > 0$  and supported in an interval of length*

$$\frac{N}{\log^{\delta-o(1)} N}$$

*which tiles with all  $\Lambda_j$ .*

*Furthermore, for any  $\epsilon > 0$  any such function  $f$  must have*

$$\text{diam supp } f \gtrsim_{\epsilon} N^{1-\epsilon}.$$

Arguing similarly we can also prove the lower bound for  $\text{diam supp } f$  in Theorem 2.9. If we assume that  $f$  tiles with all  $\Lambda_j = \lambda_j \mathbb{Z}$ , with, say,  $1 \leq \lambda_j \leq 2$ ,  $j = 1, 2, \dots, N$ , then  $\widehat{f}$  vanishes on

$$\bigcup_{j=1}^N \lambda_j^{-1} \mathbb{Z} \setminus \{0\}.$$

If this set is large then Jensen's formula implies that  $\text{diam supp } f$  is also large. It was proved in [11, Theorem 1.1, special case  $\ell = n$ ] that, for any  $\epsilon > 0$ , the above union of arithmetic progressions contains at least  $c_{\epsilon} N^{2-\epsilon}$  points in  $[0, 2N]$ . By Jensen's formula then we have  $\text{diam supp } f \gtrsim_{\epsilon} N^{1-\epsilon}$  and this completes the proof of Theorem 2.9.

**Question 6.** *Can we ensure  $f \geq 0$  in the first half of Theorem 2.9?*

## 3. THE COMMON FUNDAMENTAL DOMAIN PROBLEM WITHOUT MEASURABILITY

In [18] the following theorem was proved in §3.2.

**Theorem 3.1.** *If  $\Lambda_0, \dots, \Lambda_n$  are lattices in  $\mathbb{R}^d$  of the same volume and with the sum  $\Lambda_0 + \Lambda_1 + \dots + \Lambda_n$  being direct then there is a bounded common fundamental domain  $F$  for all these lattices.*

*Remark 3.2.* No measurability is claimed for  $F$  and the set  $F$  constructed in [18] is not measurable.

The question was left open in [18] whether the equal volume assumption was necessary. This assumption is obviously necessary if we ask for a measurable tile as the volume of the tile equals the volume of each lattice it tiles with. But there is no a priori reason for this requirement to hold if we cannot measure volumes, as with tiles that are not necessarily measurable. But, we show here that, indeed it is necessary, by giving a pair of lattices, with no common elements, which have different volumes and have no bounded common fundamental domain.

**Theorem 3.3.** *Let  $\Lambda_1 = \mathbb{Z}^d$  and  $\Lambda_2 = \alpha\mathbb{Z}^d$ , with  $\alpha$  irrational and  $d \geq 1$ . Then there is no bounded set  $F \subseteq \mathbb{R}^d$  which consists of exactly one representative from each coset of each  $\Lambda_i$ ,  $i = 1, 2$ .*

*Proof.* We give the proof in dimension  $d = 1$  for clarity, as it is essentially the same for all  $d$ . Without loss of generality we take  $\alpha > 1$ .

As explained in [18, Proof of Theorem 1] it is enough to show that  $\Lambda_1$  and  $\Lambda_2$  do not have a bounded common fundamental domain in the group

$$G = \Lambda_1 + \Lambda_2 = \{m + n\alpha : m, n \in \mathbb{Z}\}.$$

Suppose  $F$  is just such a bounded common fundamental domain

$$F = \{m_i - n_i\alpha : i = 1, 2, \dots\} \subseteq [-M, M].$$

For  $F$  to be a fundamental domain it must contain precisely one point in the set  $k + \alpha\mathbb{Z}$  and one point in the set  $\mathbb{Z} + k\alpha$ , for all  $k \in \mathbb{Z}$ . By renumbering then we can write

$$F = \{m - n_m\alpha : m \in \mathbb{Z}\}.$$

Let now  $R > 0$  be large and consider the following set of values for  $m$ :

$$(3.1) \quad -R \leq m \leq R.$$

For such values of  $m$  and from the presumed bound

$$|m - n_m\alpha| \leq M$$

we have the bounds

$$(3.2) \quad -\frac{R+M}{\alpha} \leq n_m \leq \frac{R+M}{\alpha}.$$

As  $R \rightarrow \infty$  the number of values of  $m$  allowed by (3.1) are  $\sim 2R$  in number. The number of the corresponding values of  $n$  allowed by (3.2) is  $\sim 2R/\alpha$ , which is strictly smaller if  $R$  is large since we have taken  $\alpha > 1$ . This is a contradiction as there must be exactly one  $n_m$  for each  $m$  and all the  $n_m$ 's are different.  $\square$

## 4. THE PROBLEM IN FINITE ABELIAN GROUPS

Suppose we have a finite abelian group  $G$  and two subgroups  $G_1, G_2$  of the same index

$$n = [G : G_1] = [G : G_2].$$

We can ask whether we can find a common fundamental domain for  $G_1, G_2$  in  $G$ , i.e. a set of  $n$  elements

$$F = \{g_1, \dots, g_n\}$$

which tiles with both  $G_1$  and  $G_2$ . This always exists, even in the non-abelian case, if properly defined, see e.g. [3].

If we drop the equal index assumption we can still ask for a *function*  $f$  defined on  $G$  which tiles with both subgroups:

$$(4.1) \quad \forall x \in G : \sum_{g_1 \in G_1} f(x - g_1) = |G_1|, \quad \sum_{g_2 \in G_2} f(x - g_2) = |G_2|.$$

The question we are interested in here is:

Given  $G, G_1, G_2$  how small can the size of the support of  $f$  be?

Under the assumption of equal index the answer to the above question is that, since we can find a common fundamental domain of  $G_1, G_2$  in  $G$  [18], the size of the support of  $f$  can be as small as  $[G : G_1] = [G : G_2]$ . Of course it cannot be smaller than that. But once we drop the equal index assumption then the only general construction we know is the convolution of the indicator functions of the fundamental domains  $D_i$  of  $G_i$ :

$$(4.2) \quad f = c \mathbb{1}_{D_1} * \mathbb{1}_{D_2},$$

which gives, with  $c = \frac{|G|}{|D_1| \cdot |D_2|}$ ,

$$f * \mathbb{1}_{G_1} = c \mathbb{1}_{D_2} * \mathbb{1}_{D_1} * \mathbb{1}_{G_1} = c \mathbb{1}_{D_2} * \mathbb{1}_G = c |D_2| = |G_1|,$$

and similarly for  $f * \mathbb{1}_{G_2}$ . But the support of  $f$  in (4.2) can be quite large, a priori as large as  $|D_1| \cdot |D_2|$ .

From now on we restrict our functions  $f$  to be nonnegative and normalized as shown in (4.1). We also usually restrict our tiles to be nonnegative functions.

**Definition 4.1.** If  $G_1, G_2$  are subgroups of the finite abelian group  $G$  we write

$$S_{G_1, G_2}^G = \min \left\{ |\text{supp } f| : \text{where } f : G \rightarrow \mathbb{R}^{\geq 0}, f * \mathbb{1}_{G_1} = |G_1|, f * \mathbb{1}_{G_2} = |G_2| \right\}.$$

It is always the case that

$$S_{G_1, G_2}^G \geq [G : G_i], \quad (i = 1, 2).$$

Observe that if  $G_1, G_2$  have a common fundamental domain  $F$  in  $G$  then

$$S_{G_1, G_2}^G = |F|.$$

The following result says that we can always restrict our study to the case of  $G_1 \cap G_2$  being trivial. In other words we may assume from now on that  $G$  is the direct sum of  $G_1$  and  $G_2$ .

**Theorem 4.2.** If  $G_1, G_2 \subseteq G$  are finite abelian groups and

$$\Gamma = G / (G_1 \cap G_2), \quad \Gamma_i = G_i / (G_1 \cap G_2), \quad (i = 1, 2)$$

then

$$(4.3) \quad S_{G_1, G_2}^G = S_{\Gamma_1, \Gamma_2}^\Gamma.$$

*Proof.* In what follows  $\Gamma$  or  $\Gamma_i$  may denote either the quotient group or an arbitrary but fixed set of coset representatives of the subgroup  $G_1 \cap G_2$ .

Let  $f : G \rightarrow \mathbb{R}^{\geq 0}$  satisfy

$$|G_i| = f * \mathbb{1}_{G_i} = (f * \mathbb{1}_{G_1 \cap G_2}) * \mathbb{1}_{\Gamma_i}, \quad (i = 1, 2).$$

If  $\text{supp } f$  is minimal it follows from the above representation that  $\text{supp } f$  has at most one point in each  $(G_1 \cap G_2)$ -coset, as we can always collect all the ‘‘mass’’ of  $f$  contained in one coset to one point of the coset without affecting  $f * \mathbb{1}_{G_1 \cap G_2}$ .

Define  $F : \Gamma \rightarrow \mathbb{R}^{\geq 0}$  by

$$F(\gamma) = \frac{1}{|G_1 \cap G_2|} \sum_{g \in G_1 \cap G_2} f(\gamma + g) = \frac{1}{|G_1 \cap G_2|} f * \mathbb{1}_{G_1 \cap G_2}(\gamma).$$

It follows, under the assumption that  $|\text{supp } f|$  is minimal, that

$$|\text{supp } F| = |\text{supp } f|.$$

We also have, for  $i = 1, 2$ ,

$$F * \mathbb{1}_{\Gamma_i} = \frac{1}{|G_1 \cap G_2|} f * \mathbb{1}_{G_1 \cap G_2} * \mathbb{1}_{\Gamma_i} = \frac{1}{|G_1 \cap G_2|} f * \mathbb{1}_{G_i} = \frac{|G_i|}{|G_1 \cap G_2|} = |\Gamma_i|.$$

It follows that

$$S_{\Gamma_1, \Gamma_2}^\Gamma \leq S_{G_1, G_2}^G.$$

To prove the reverse inequality we start with a function  $F : \Gamma \rightarrow \mathbb{R}^{\geq 0}$  satisfying

$$F * \mathbb{1}_{\Gamma_i} = |\Gamma_i|, \quad (i = 1, 2)$$

and define  $f : G \rightarrow \mathbb{R}^{\geq 0}$  by taking  $f(x)$  to be

$$|G_1 \cap G_2| \cdot F(x + G_1 \cap G_2)$$

at precisely one point  $x$  in each  $(G_1 \cap G_2)$ -coset, and in all other points of the coset we take it to be 0.

It follows that  $|\text{supp } f| = |\text{supp } F|$ , and we also have, for  $i = 1, 2$ , and viewing  $F$  as a function on  $G$  (constant on  $(G_1 \cap G_2)$ -cosets),

$$f * \mathbb{1}_{G_i} = f * \mathbb{1}_{G_1 \cap G_2} * \mathbb{1}_{\Gamma_i} = |G_1 \cap G_2| \cdot F * \mathbb{1}_{\Gamma_i} = |G_1 \cap G_2| \cdot |\Gamma_i| = |G_i|.$$

This concludes the proof of the reverse inequality and the Theorem. □

Let  $G = G_1 \times G_2$  from now on. The following theorem is the best understood case.

**Theorem 4.3.** *If  $G = G_1 \times G_2$  and  $|G_1|$  divides  $|G_2|$  then*

$$(4.4) \quad S_{G_1, G_2}^G = [G : G_1] = |G_2|.$$

*Proof.* Enumerate the two subgroups arbitrarily as

$$G_1 = \{g_1^1, \dots, g_m^1\}, \quad G_2 = \{g_1^2, \dots, g_n^2\},$$

with  $n = km$ ,  $k \geq 1$ . Take  $f = |G_1| \mathbb{1}_F$  where the set  $F$  is constructed by taking sums of the elements of  $G_2$  with the ‘‘corresponding’’ elements of  $G_1$  (but the

elements of  $G_1$  have to be repeated  $k$  times each). So  $F$  consists of the sums

$$\begin{aligned} &g_1^1 + g_1^2, g_2^1 + g_2^2, \dots, g_m^1 + g_m^2 \\ &g_1^1 + g_{m+1}^2, g_2^1 + g_{m+2}^2, \dots, g_m^1 + g_{m+m}^2 \\ &g_1^1 + g_{2m+1}^2, \dots, g_m^1 + g_{2m+m}^2 \\ &\dots \end{aligned}$$

It is easy to see that  $F$  is a fundamental domain for  $G_1$  and that

$$f * \mathbb{1}_{G_1} = |G_1|, \quad f * \mathbb{1}_{G_2} = k|G_1| = |G_2|.$$

Finally  $|\text{supp } f| = |G_2|$ , which proves the Theorem as we always have  $S_{G_1, G_2}^G \geq |G_2|$ .  $\square$

It is clear now that in studying the problem in the group  $G = G_1 \times G_2$  the group structure is irrelevant and, writing  $m = |G_1|$  and  $n = |G_2|$ , the problem is to find a nonnegative real matrix

$$A \in (\mathbb{R}^{\geq 0})^{m \times n}$$

with row sums all equal to  $n$  and column sums all equal to  $m$ , and with as small a support (non-zero entries) as possible.

Matrices of this type or, rather, the matrices  $\frac{1}{mn}A$ , and, more generally, multivariate distributions with uniform marginals, are called *copulas* in statistics and have been studied extensively [27]. They can be used to “isolate” the marginal distributions of a general multivariate distribution from the dependence part of distribution.

**Definition 4.4.** Write  $A(m, n)$  for the set of all such  $m \times n$  matrices (with non-negative entries and all row sums equal to  $n$  and all column sums equal to  $m$ ) and

$$S(m, n) = \min \left\{ |\text{supp } A| : A \in A(m, n) \right\}.$$

In this notation Theorem 4.3 says that

$$S(m, km) = km, \quad \text{if } m, k \in \mathbb{N}.$$

In what follows we use [4], where the structure of the matrices in  $A(m, n)$  of minimal support is described up to permutation of rows and columns (these operations obviously leave  $A(m, n)$  unchanged and also do not alter the size of the support of each matrix).

That the situation changes radically if  $m$  does not divide  $n$  can be seen, for example, by the following.

**Lemma 4.5.** *If  $k \geq 1$ ,  $1 \leq r < m$ , then*

$$S(m, km + r) \geq (k + 1)m.$$

*If  $r = 1$  we have*

$$S(m, km + 1) = (k + 1)m.$$

*Proof.* Column sums are equal to  $m$ , so none of the entries is  $> m$ . Since row sums are equal to  $km + r$  it follows that we have at least  $k + 1$  non-zero terms

in each of the  $m$  rows. This shows the lower bound  $S(m, km + r) \geq m(k + 1)$ . To show the next claim we check that the  $m \times (km + 1)$  matrix

$$m \left\{ \begin{array}{cccc} & \overbrace{m \cdots m}^k & \cdots & \cdots \\ 1 & & \overbrace{m \cdots m}^k & \cdots \\ & \cdots & & \cdots \\ & \cdots & & \overbrace{m \cdots m}^k \\ 1 & \cdots & \cdots & \end{array} \right\}$$

$km+1$

has constant row and column sums. Since its support has size  $(k + 1)m$ , which is the minimum possible by the first part of the Lemma, we are done.  $\square$

**Question 7.** *What is the true value of  $S(m, km + r)$  when  $1 < r < m$ ?*

The construction given in Lemma 4.5 can be transferred to the real line as follows.

**Theorem 4.6.** *If  $k \geq 1$ ,  $1 \leq r < m$ , and  $\Lambda_1 = m\mathbb{Z}$ ,  $\Lambda_2 = (km + 1)\mathbb{Z}$  then there exists a measurable  $F : \mathbb{R} \rightarrow \mathbb{R}^{\geq 0}$  which tiles  $\mathbb{R}$  with both  $\Lambda_1$  and  $\Lambda_2$  and has*

$$|\text{supp } F| = (k + 1)m.$$

*Remark 4.7.* Notice that  $(k + 1)m$  is 1 less than the size of the support that we would get had we used the convolution (2.7)  $\mathbb{1}_{[0,m]} * \mathbb{1}_{[0,km+1]}$ . Therefore the convolution tile is not optimal in terms of its length, at least for the lattices  $m\mathbb{Z}$  and  $(km + 1)\mathbb{Z}$ .

*Proof.* We can view the function  $f$  constructed in the second part of Lemma 4.5 as a function on the group  $\mathbb{Z}_m \times \mathbb{Z}_n$ , with  $n = km + 1$ . Since  $m$  and  $km + 1$  are coprime this group is isomorphic to the cyclic group  $\mathbb{Z}_{mn}$  and we can view our function  $f$  as being defined on  $\mathbb{Z}_{mn}$  and tiling with both subgroups  $\langle m \rangle$  and  $\langle n \rangle$ , of size  $n$  and  $m$  respectively, therein (here  $\langle g \rangle$  denotes the subgroup generated by  $g$ ):

$$\forall x \in \mathbb{Z}_{mn} : \sum_{g \in \langle n \rangle} f(x - g) = m, \quad \sum_{g \in \langle m \rangle} f(x - g) = n.$$

This is the same as tiling  $\mathbb{Z}$  (now  $\langle n \rangle = n\mathbb{Z}$ ,  $\langle m \rangle = m\mathbb{Z}$ )

$$\forall x \in \mathbb{Z} : \sum_{g \in \langle n \rangle} f(x - g) = m, \quad \sum_{g \in \langle m \rangle} f(x - g) = n.$$

Finally, defining  $F = f * \mathbb{1}_{[0,1]}$ , we obtain tilings of  $\mathbb{R}$ :

$$\forall x \in \mathbb{R} : \sum_{g \in \langle n \rangle} F(x - g) = m, \quad \sum_{g \in \langle m \rangle} F(x - g) = n.$$

All that remains is to observe that  $|\text{supp } F| = (k + 1)m$ .

$\square$

## REFERENCES

1. J. Beck, *On a lattice point problem of H. Steinhaus*, *Studia Sci. Math. Hung* **24** (1989), 263–268. [2](#), [3](#)
2. A Beurling, *Collected Works of Arne Beurling (2 vol.)*, edited by L. Carleson et al, 1989. [9](#)
3. J. Button, M. Chiodo, and M. Zeron-Medina Laris, *Coset intersection graphs for groups*, *The American Mathematical Monthly* **121** (2014), no. 10, 922–926. [11](#)
4. R. M. Caron, X. Li, P. Mikusiński, H. Sherwood, and M. D. Taylor, *Nonsquare “Doubly Stochastic” Matrices*, *Lecture Notes-Monograph Series* (1996), 65–75. [13](#)
5. W. K. Chan and R. Mauldin, *Steinhaus tiling problem and integral quadratic forms*, *Proceedings of the American Mathematical Society* **135** (2007), no. 2, 337–342. [2](#)
6. M. Ciucu, *A remark on sets having the Steinhaus property*, *Combinatorica* **16** (1996), no. 3, 321–324. [2](#)
7. H. T. Croft, *Three lattice-point problems of Steinhaus*, *The Quarterly Journal of Mathematics* **33** (1982), no. 1, 71–83. [2](#)
8. W. F. Donoghue, *Distributions and Fourier transforms*, Academic Press, 2014. [4](#)
9. Paul Erdos, *Note on sequences of integers no one of which is divisible by any other*, *J. London Math. Soc* **10** (1935), no. 1, 126–128. [9](#)
10. R. Gardner, *The Brunn-Minkowski inequality*, *Bulletin of the American Mathematical Society* **39** (2002), no. 3, 355–405. [6](#)
11. S. Gilboa and R. Pinchasi, *On the union of arithmetic progressions*, *SIAM Journal on Discrete Mathematics* **28** (2014), no. 3, 1062–1073. [9](#)
12. V. Havin and B. Jöricke, *The Uncertainty Principle in Harmonic Analysis*, Springer, Berlin, 1994. [2](#)
13. S. Jackson and R. Mauldin, *On a lattice problem of H. Steinhaus*, *Journal of the American Mathematical Society* **15** (2002), no. 4, 817–856. [2](#)
14. S. Jackson and R. D. Mauldin, *Sets meeting isometric copies of the lattice  $\mathbb{Z}^2$  in exactly one point*, *Proceedings of the National Academy of Sciences* **99** (2002), no. 25, 15883–15887. [2](#)
15. ———, *Survey of the Steinhaus tiling problem*, *Bulletin of Symbolic Logic* **9** (2003), no. 03, 335–361. [2](#)
16. M. N. Kolountzakis, *A new estimate for a problem of Steinhaus*, *International Mathematics Research Notices* **1996** (1996), no. 11, 547–555. [2](#), [3](#)
17. ———, *A problem of Steinhaus: Can all placements of a planar set contain exactly one lattice point?*, *Progress in Mathematics* **139** (1996), 559–566. [2](#), [3](#)
18. ———, *Multi-lattice tiles*, *International Mathematics Research Notices* **1997** (1997), no. 19, 937–952. [2](#), [3](#), [7](#), [10](#), [11](#)
19. ———, *The study of translational tiling with Fourier Analysis*, *Fourier Analysis and Convexity* (L. Brandolini, ed.), Birkhäuser, 2004, pp. 131–187. [2](#), [4](#)
20. M. N. Kolountzakis and M. Papadimitrakis, *The Steinhaus tiling problem and the range of certain quadratic forms*, *Illinois Journal of Mathematics* **46** (2002), no. 3, 947–951. [2](#)
21. ———, *Measurable steinhaus sets do not exist for finite sets or the integers in the plane*, *Bulletin of the London Mathematical Society* **49** (2017), no. 5, 798–805. [2](#)
22. M. N. Kolountzakis and T. Wolff, *On the Steinhaus tiling problem*, *Mathematika* **46** (1999), no. 02, 253–280. [2](#), [3](#), [4](#), [5](#)
23. P. Komjáth, *A lattice-point problem of Steinhaus*, *The Quarterly Journal of Mathematics* **43** (1992), no. 2, 235–241. [2](#)
24. E. Mallinikova and S. Rukshin, *On one Steinhaus problem*, *Vestnik St.Petersburg Univ. Math.* **28** (1995), no. 1, 28–32. [2](#)
25. R. Mauldin and A. Yingst, *Comments about the Steinhaus tiling problem*, *Proceedings of the American Mathematical Society* **131** (2003), no. 7, 2071–2079. [2](#)
26. R. D. Mauldin, *Some problems in set theory, analysis and geometry*, *Paul Erdos and his Mathematics I* (2001), 493–505. [2](#)
27. R. B. Nelsen, *An introduction to copulas*, Springer Science & Business Media, 2007. [13](#)
28. W. Sierpiński, *Sur un probleme de H. Steinhaus concernant les ensembles de points sur le plan*, *Fundamenta Mathematicae* **2** (1958), no. 46, 191–194. [2](#)
29. S. M. Srivastava and R. Thangadurai, *On Steinhaus sets*, *Expositiones Mathematicae* **23** (2005), no. 2, 171–177. [2](#)
30. G. Tenenbaum, *Lois de répartition des diviseurs*, *Séminaire Delange-Pisot-Poitou. Théorie des nombres* **19** (1980), no. 1, 1–3. [9](#)

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, UNIVERSITY OF CRETE, VOUTES CAMPUS, 70013 HERAKLION, CRETE, GREECE.

*Email address:* kolount@gmail.com

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, UNIVERSITY OF CRETE, VOUTES CAMPUS, 70013 HERAKLION, CRETE, GREECE.

*Email address:* papageoeffie@gmail.com