

# A remark on perturbations of sine and cosine sums

MIHAIL N. KOLOUNTZAKIS

December 1999

Consider a collection  $\lambda_1 < \dots < \lambda_N$  of distinct positive integers and the quantities

$$M_1 = M_1(\lambda_1, \dots, \lambda_N) = \max_{0 \leq x \leq 2\pi} \left| \sum_{j=1}^N \sin \lambda_j x \right|$$

and

$$M_2 = M_2(\lambda_1, \dots, \lambda_N) = - \min_{0 \leq x \leq 2\pi} \sum_{j=1}^N \cos \lambda_j x.$$

One is interested in frequencies  $\lambda_j$  which make the quantities  $M_1$  and  $M_2$  small as  $N \rightarrow \infty$ . It is trivial that  $M_1 \geq cN^{1/2}$  but it is much harder to show even that  $M_2 \rightarrow \infty$ . (In this note  $c$  denotes an absolute positive constant, not necessarily the same in all its occurrences.)

It is a result of Bourgain [1, 2] that  $M_1$  may become  $O(N^{2/3})$  and it is very easy to construct a collection  $\lambda_j$  which gives  $M_2 = O(N^{1/2})$ , which is the conjectured optimal. For minimizing  $M_2$  it is also possible to have the collection of frequencies relatively well packed, that is with  $\lambda_N \leq 2N$  [3], while for any  $\epsilon > 0$  and for any collection  $\lambda_j$  that makes  $M_1 = O(N^{1-\epsilon})$  one can easily see that  $\lambda_N$  is super-polynomial in  $N$ .

Prompted by a discussion with G. Benke we prove that collections of frequencies  $\lambda_j$  which have  $M_1 = o(N)$  or  $M_2 = o(N)$  are unstable, in the sense that one can perturb the  $\lambda_j$  by one each and get  $M_1 \geq cN$  and  $M_2 \geq cN$ .

**Theorem 1** (i) *Suppose that  $M_1(\lambda_1, \dots, \lambda_N) = o(N)$ . Then there exists a choice of  $\epsilon_j = \pm 1$ ,  $j = 1, \dots, N$ , so that  $M_1(\lambda_1 + \epsilon_1, \dots, \lambda_N + \epsilon_N) \geq cN$ .*  
(ii) *Suppose, similarly, that  $M_2(\lambda_1, \dots, \lambda_N) = o(N)$ . Then there exists a choice of  $\epsilon_j = \pm 1$ ,  $j = 1, \dots, N$ , so that  $M_2(\lambda_1 + \epsilon_1, \dots, \lambda_N + \epsilon_N) \geq cN$ .*

**Remark.** It is not always the case that the perturbed frequencies are all distinct but is frequently so and, in any case, at most two may overlap at any given integer.

**Proof.** (i) Write

$$\begin{aligned} \sum_{j=1}^N \sin(\lambda_j + \epsilon_j)x &= \sum_{j=1}^N \sin \lambda_j x \cos \epsilon_j x + \sum_{j=1}^N \cos \lambda_j x \sin \epsilon_j x \\ &= \cos x \sum_{j=1}^N \sin \lambda_j x + \sin x \sum_{j=1}^N \epsilon_j \cos \lambda_j x \\ &= \text{I} + \text{II}. \end{aligned}$$

We have  $\text{I} = o(N)$ . For  $\lambda > 10$ , say, we have

$$\frac{4}{\pi} \int_{\pi/4}^{\pi/2} |\cos \lambda x| dx \geq c.$$

From this we deduce that

$$\frac{4}{\pi} \int_{\pi/4}^{\pi/2} \sum_{j=1}^N |\cos \lambda_j x| dx \geq cN,$$

hence there exists  $x_0 \in [\frac{\pi}{4}, \frac{\pi}{2}]$  such that

$$\sum_{j=1}^N |\cos \lambda_j x_0| \geq cN.$$

Choose then  $\epsilon_j = \text{sgn}(\cos \lambda_j x_0)$  to get

$$\sum_{j=1}^N \epsilon_j \cos \lambda_j x_0 \geq cN.$$

Since  $\sin x \geq 2^{-1/2}$  in  $[\frac{\pi}{4}, \frac{\pi}{2}]$  we get that  $\text{II} \geq cN$  at  $x_0$ , which gives the required  $M_1(\lambda_1 + \epsilon_1, \dots, \lambda_N + \epsilon_N) \geq cN$ , since  $\text{I} = o(N)$  everywhere.

(ii) The proof is similar. We write

$$\begin{aligned} \sum_{j=1}^N \cos(\lambda_j + \epsilon_j)x &= \cos x \sum_{j=1}^N \cos \lambda_j x - \sin x \sum_{j=1}^N \epsilon_j \sin \lambda_j x \\ &= \text{I} - \text{II}. \end{aligned}$$

For  $x \in [\frac{4\pi}{6}, \frac{5\pi}{6}]$  we have  $\text{I} \leq o(N)$  so it is enough to show that for some  $x_0$  in the same interval  $\text{II} \geq cN$ . To do that observe, as before, that

$$\frac{6}{\pi} \int_{4\pi/6}^{5\pi/6} \sum_{j=1}^N |\sin \lambda_j x| \geq cN, \quad (1)$$

and choose  $\epsilon_j = \text{sgn}(\sin \lambda_j x_0)$ , where  $x_0$  is a point that makes the left hand side of (1) large. This implies the existence of  $x_0 \in [\frac{4\pi}{6}, \frac{5\pi}{6}]$  such that  $\text{II} \geq cN$  at  $x_0$ .

□

Take a Bourgain's collection of frequencies  $\lambda_j$  for which  $M_1 = O(N^{2/3})$ . Bourgain gave a randomized construction for these which produces  $N$  frequencies in the interval  $[1, e^{N^{1/3}}]$ . By taking the inner product of the above sine sum with the conjugate Dirichlet kernel  $D_M^* = \sum_{j=1}^M \sin jx$ , whose  $L^1$  norm is  $c \log M$ , we obtain that the number of frequencies in the interval  $[1, M]$  is at most  $cN^{2/3} \log M$ , hence Bourgain's method does not go any further than it has to (in the size of the  $\lambda_j$ ). One can even modify his randomized construction to produce  $\lambda_j \sim e^{j^{1/3}}$ . Hence the question becomes reasonable of whether it is just the growth of the frequencies that achieves the result. Our Theorem 1 answers this in the negative.

## Bibliography

- [1] J. Bourgain, Sur les sommes de sinus, Sém. Anal. Harm., Publ. Math. d'Orsay 84-01 (1984), exp. no 3.
- [2] J.-P. Kahane, *Some random series of functions*, Cambridge Studies in Advanced Mathematics 5, 1985, Second Edition.
- [3] M.N. Kolountzakis, On nonnegative cosine polynomials with nonnegative, integral coefficients, Proc. Amer. Math. Soc. **120** (1994), vol. 1, 157-163.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CRETE, KNOSSOS AVE., 714 09 IRAKLIO, GREECE.

E-MAIL: kolount@math.ucl.ac.uk